

## ABSTRACT\*

### **Selective stimulation of the growth of anaerobic microflora in the human intestinal tract by electrolyzed reducing water.**

Department of Physiology of Microorganisms, Biology Faculty, Lomonosov Moscow State University, 119992 Moscow, Russia. Vorobjeva NV, Med Hypotheses. 2005;64(3):543 -6.

**96-99% of the "friendly" or residential microflora of intestinal tract of humans consists of strict anaerobes and only 1-4% of aerobes. Many diseases of the intestine are due to a disturbance in the balance of the microorganisms inhabiting the gut.** The treatment of such diseases involves the restoration of the quantity and/or balance of residential microflora in the intestinal tract. It is known that aerobes and anaerobes grow at different oxidation-reduction potentials (ORP). The former require positive E(h) values up to +400 mV. Anaerobes do not grow unless the E(h) value is negative between -300 and -400 mV. In this work, it is suggested that prerequisite for the recovery and maintenance of obligatory anaerobic microflora in the intestinal tract is a negative ORP value of the intestinal milieu. **Electrolyzed reducing water with E(h) values between 0 and -300 mV produced in electrolysis devices possesses this property. Drinking such water favours the growth of residential microflora in the gut. A sufficient array of data confirms this idea. However, most researchers explain the mechanism of its action by an antioxidant properties destined to detox the oxidants in the gut and other host tissues. Evidence is presented in favour of the hypothesis that the primary target for electrolyzed reducing water is the residential microflora in the gut.**

PMID: 15617863 [PubMed - indexed for MEDLINE]

\*In regard to this study, we only have the abstract and do not own the full study. The abstract explains the brief results and outcome of the study that was performed.